

CHARACTERS OF REPRESENTATIONS OF QUANTUM GROUPS OF TYPE A_n

PHÙNG HỒ HẢI

ABSTRACT. We introduce the notion of characters of comodules over coribbon Hopf algebras. The case of quantum groups of type A_n is studied. We establish a characteristic equation for the quantum matrix and a q -analogue of Harish-Chandra-Itzykson-Zuber integral

INTRODUCTION

Let G be an algebraic group, V be a G -module. The character Φ_V of V is a function on G , whose value at an element g of G is the trace of the operator on V , induced by g . In this note we introduce the notion of characters of quantum groups. Thus the algebraic group G will be replaced by a coribbon Hopf algebra H , G -module will be replaced by H -comodules, their characters are elements of H .

To the further generalization, we also introduce the notion of characters of endomorphisms of H -comodules, thus obtain a trace map from the endomorphism rings of H -comodules into H . Further we introduce the notion of partial characters. As an application we define the braided powers of a multiplicative matrix.

We focus ourselves on matrix quantum groups of type A_{n-1} . These quantum groups are defined in terms of even Hecke symmetries. They are quantum analogies of the general linear group $GL(n)$, whose characters are closely related to symmetric functions in n variables. It turns out that the characters of a quantum group of type A_{n-1} form a ring that is isomorphic to the ring of symmetric functions in n variables. The characters of endomorphisms of the tensor powers of the basic comodule induces a trace map on the Hecke algebra with values in the ring of characters. Finally, using the partial characters we define the quantum powers of a quantum matrix and prove the characteristic equation for these powers.

As an application, we give a q -analogue of the Harish-Chandra-Itzykson-Zuber (HCIZ) integral formula. Classically, the HCIZ formula computes the integral $I(M, N, t) := \int_{U(n)} \exp(\text{tr}(MUNU^\dagger)) dU$ on the unitary group $U(n)$, where M, N are hermitian matrices. This integral is computed in terms of irreducible characters of $U(n)$ evaluated at M and N [1]:

$$I(M, N, t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{\lambda \vdash n} \frac{d_\lambda}{r_\lambda} \Phi_\lambda(M) \Phi_\lambda(N).$$

We also derive a formula for a special trace map of the Hecke algebras at primitive idempotents. Consequently, we obtained an interesting interpretations of the characters tables of the Hecke algebras.

The paper is organized as follows. In Section 1 we give the definitions and basic properties of coquasitriangular and coribbon Hopf algebras. Then we define the characters of comodules over coribbon Hopf algebras as well as characters of endomorphisms and partial characters. In Section 2 we first recall the definitions and representation theory of a matrix quantum group of type A_{r-1} . It is defined

in terms of an even Hecke symmetry R of rank r over a vector space V . For more details the reader is referred to [4, 5]. The Hecke symmetry induces an action of the Hecke algebra over the tensor powers of V , which is the centralizer of the action of the quantum group on V . The character map then induces a trace map on the Hecke algebras with values in the commutative algebra spanned by characters. We then define the quantum powers of the quantum matrix and prove the characteristic equation. In Section 3 we prove a quantum analogue to the HCIZ formula for the integral on quantum groups of type A_{r-1} in terms of the just introduced characters. At the end we also notice an interesting interpretation of the characters tables of Hecke algebras in terms of their minimal central idempotents.

Notations. All bialgebras, Hopf algebras as well as their comodules are defined over a fixed algebraically closed field $\mathbb{K}$ of characteristic zero. For a bialgebra, the coproduct is denoted by Δ , the counit is denoted by ε . For a Hopf algebra, the antipode is denoted by S .

The Hecke algebra $\mathcal{H}_n = \mathcal{H}_{q,n}$ is spanned as a vector space over $\mathbb{K}$ by a basis $\{T_w | w \in \mathfrak{S}_n\}$, $\mathfrak{S}_n$ is the permutation group. The element $T_{(i,i+1)}$ is denoted by T_i . $\mathcal{H}_n$ is generated, as an algebra, by $\{T_i, i = 1, 2, \dots, n-1\}$ subject to the relation $T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}$, $T_i^2 = (q-1)T_i + q$. It is semisimple provided $q^n \neq 1, \forall n > 1$.

A partition λ of a positive integers n is a sequence of non-increasing non-negative integers, whose sum is n , in notation $\lambda \vdash n$ or $|\lambda| = n$. $\mathcal{P}$ denotes the set of all partitions, $\mathcal{P}_n$ denotes the set of partitions of an integer n , $\mathcal{P}^r$ denotes the set of all partitions of length (i.e., the number of non-zero components) at most r .

A diagram $[\lambda]$ is a matrix, whose first row contains λ_1 entries, second row contains λ_2 entries, and so on... The coordinate of a node in the i -th row and j -th column is (i, j) .

1. CORIBBON HOPF ALGEBRAS AND THEIR CHARACTERS

1.1. Definitions. A coquasitriangular (CQT) structure on a bialgebra B is a bilinear form $r : B \otimes B \rightarrow \mathbb{K}$ subject to the following conditions

- i. $a_{(1)}b_{(1)}r(a_{(2)}, b_{(2)}) = r(a_{(1)}, b_{(1)})b_{(2)}a_{(2)} - \text{“naturalness”}$,
- ii. r is invertible in $\text{Hom}(B \otimes B, \mathbb{K})$,
- iii. $r(a, bc) = r(a_{(2)}, b)r(a_{(1)}, c)$
 $r(ab, c) = r(a, c_{(1)})r(b, c_{(2)}) - \text{“multiplicativity”}$.

Here we use Sweedler's sigma notation for the coproduct: $\Delta(x) = x_{(1)} \otimes x_{(2)}$.

A (Hopf) bialgebra equipped with such a structure is called coquasitriangular (CQT) (Hopf) bialgebra.

In a CQT Hopf algebra, the square of the antipode is co-inner, that is

$$S^2(a) = u^{-1}(a_{(1)})a_{(2)}u(a_{(3)}), \quad (1)$$

where $u(a) := r(a_{(2)}, S(a_{(1)}))$ and u^{-1} is its inverse in $\text{Hom}(B, \mathbb{K})$, $u^{-1}(a) = r(S^2(a_{(2)}), a_{(1)})$.

A CQT structure on a bialgebra induces a braiding [6] in the category of its right comodules in the following way. Let (B, r) be a CQT bialgebra, for any right B -comodules M, N , define a morphism

$$\tau_{M,N}(m, n) = n_{(0)} \otimes m_{(0)} r(m_{(1)}, n_{(1)}) : M \otimes N \rightarrow N \otimes M.$$

Then τ is a braiding in $B\text{-Comod}$ – the category of right B -comodules.

A coribbon Hopf algebra is a CQT Hopf algebra (H, r) equipped with a linear mapping $t : B \rightarrow \mathbb{K}$ subject to the following relations:

- i. $t(a_{(1)})a_{(2)} = a_{(1)}t(a_{(2)}) - \text{“naturalness”}$,
- ii. $t \circ S = t - \text{“rigidity”}$,
- iii. t is invertible in $\text{Hom}(B, \mathbb{K})$,
- iv. $t(ab) = t(a_{(1)})t(b_{(1)})r(b_{(2)}a_{(2)})r(a_{(3)}, b_{(3)}) - \text{“twist”}$.
- v. $t(1_H) = 1 - \text{“normalization”}$.

t has the following property:

$$t(a_{(1)})t(a_{(2)}) = u^{-1}(a_{(1)})u^{-1}(S(a_{(2)})). \quad (2)$$

The proof is to write out the right-hand side of the quality $\varepsilon(a) = t(a_{(1)})S(a_{(2)})$ and to use (1).

Let (H, r, t) be a coribbon Hopf algebra. Then H -comod, the category of finite dimensional right H -comodules, is a ribbon category [9, 6] with the twist given by

$$\theta_M : M \longrightarrow M; \quad \theta_M(m) = m_{(0)}t(m_{(1)}), \quad (3)$$

for any finite dimension right H -comodule M . Then θ commutes with any morphism and we have

$$\begin{aligned} \theta_{M^*} &= \theta_M^* & \theta_{\mathbb{K}} &= \text{id}_{\mathbb{K}} \\ \theta_{M \otimes N} &= (\theta_M \otimes \theta_N)\tau_{N,M}\tau_{M,N}. \end{aligned}$$

1.2. The Characters. Now we are going to define the characters of comodules over a coribbon Hopf algebra. First we recall the definition of rank or braided dimension of comodules. The rank of a comodule M over a coribbon Hopf algebra H is

$$\text{rank}(M) := \text{ev}_V \circ \tau_{M,M^*}(\theta_M \otimes \text{id}_{M^*})\text{db}_M(1_{\mathbb{K}}) \in \mathbb{K},$$

where $\text{db}_M : \mathbb{K} \longrightarrow M \otimes M^*$, $\text{ev}_M : M^* \otimes M \longrightarrow \mathbb{K}$ are morphisms that make M^* the dual comodule to M . The isomorphism θ is inserted in the definition in order to provide that $\text{rank}(M \otimes N) = \text{rank}(M) \cdot \text{rank}(N)$. It is also obvious that $\text{rank}(M \oplus N) = \text{rank}(M) + \text{rank}(N)$, $\text{rank}(M^*) = \text{rank}(M)$. Unfortunately, the rank is generally not additive with respect to short exact sequences. For a morphism $f : M \longrightarrow M$, we define its braided trace to be

$$\text{Tr}(f) := \text{ev}_V \circ \tau_{M,M^*}(f \circ \theta_M \otimes \text{id}_{M^*})\text{db}_M(1_{\mathbb{K}}) \in \mathbb{K}.$$

It is easy to see that Tr is additive and tensor multiplicative, moreover, $\text{Tr}(f \circ g) = \text{Tr}(g \circ f)$. Notice that $\text{rank}(M) = \text{Tr}(\text{id}_M)$.

The character $\Phi(M)$ is defined to be

$$\Phi(M) := (\text{ev}_M \otimes \text{id}_H)(\text{id}_{M^*} \otimes \delta_M)\tau_{M,M^*}(\theta_M \otimes \text{id}_{M^*})\text{db}_M(1_{\mathbb{K}}) \in H.$$

Analogously, for a morphism $f : M \longrightarrow M$, we define its character to be

$$\Phi(f) := (\text{ev}_M \otimes \text{id}_H)(\text{id}_{M^*} \otimes \delta_M)\tau_{M,M^*}(f \circ \theta_M \otimes \text{id}_{M^*})\text{db}_M(1_{\mathbb{K}}) \in H.$$

Notice that $\Phi(M) = \Phi(\text{id}_M)$ and $\varepsilon(\Phi(f)) = \text{Tr}(f)$, ε is the counit on H . Thus we have a character map $\Phi : \text{End}^H(M) \longrightarrow H$.

Lemma 1. *The character map Φ has the following properties*

- i. $\Phi(f + g) = \Phi(f) + \Phi(g)$
- ii. $\Phi(f \otimes h) = \Phi(f)\Phi(h)$
- iii. $\Phi(f \circ g) = \Phi(g \circ f)$.

(ii) and (iii) imply that $\Phi(f)\Phi(g) = \Phi(g)\Phi(f)$. *Proof.* . The properties (i) and (iii) are obviously. (ii) follows from the following identity

$$\tau_{M^{\otimes n}, M^{*\otimes n}}(\theta_{M^{\otimes n}} \otimes \text{id}_{M^{*\otimes n}}) = \text{db}_M^n \circ \text{db}_M^{n-1} \circ \dots \circ \text{db}_M : \mathbb{K} \longrightarrow M^{\otimes n} \otimes M^{*\otimes n}, \quad (4)$$

where $\text{db}_{M^{\otimes n}}^i := \text{id}_M^{i-1} \otimes \text{db}_M \otimes \text{id}_{M^*}^{i-1}$.

Thus, Φ is a trace map on $\text{End}^H(M)$ with value in the commutative subalgebra of H spanned by characters.

Assume now that $f : M \otimes N \longrightarrow M \otimes N$. Then we define its partial character with respect to M , $\Phi_M(f) : N^* \otimes N \longrightarrow H$, to be

$$\Phi_M(f) := (\text{ev}_{M \otimes N} \otimes \text{id}_H)\delta_{M \otimes N}(\text{id}_{N^*} \otimes (f \circ \tau_{M,M^*}(\theta_M \otimes \text{id}_{M^*})\text{db}_M) \otimes \text{id}_N).$$

The partial character is used particularly to define the braided powers of the multiplicative matrix associated to M . We mention the following property of the partial

character, which will be used later. Let $f : M \longrightarrow M$ and $g : M \otimes N \longrightarrow M \otimes N$. Then

$$\Phi_M((f \otimes \text{id}_N) \circ g) = \Phi_M(g \circ (f \otimes \text{id}_N)). \quad (5)$$

Let us now give some coordinate calculation. Fix a basis $x_1, x_2, \dots, x_n$ of M . The coaction of H can then be given in terms of a multiplicative matrix $A = (a_j^i)_{i,j=1}^n$: $\delta(x_i) = x_j \otimes a_i^j$, $\Delta(a_j^i) = a_k^i \otimes a_j^k$, here and later on we shall adopt the convention of summing up by the indices that appear both in upper and lower places. Let $\xi_1, \xi_2, \dots, \xi_n$ be the dual basis on M^* . Thus, $\text{db}_M(1_{\mathbb{K}}) = x_i \otimes \xi^i$, $\text{ev}_M(\xi^i \otimes x_j) = \delta_j^i$.

Assume that with respect to the bases above $\tau_{M,M}$ and θ_M have matrices R_{kl}^{ij} and T_j^i , respectively:

$$\tau_{M,M}(x_i \otimes x_j) = x_k \otimes x_l R_{ij}^{kl}, \quad \theta_M(x_i) = x_j T_i^j.$$

Then we have $r(a_j^i, a_l^k) = R_{jl}^{ki}$, $t(a_j^i) = T_j^i$. Set $P_{jl}^{ki} = r(a_j^i, S(a_l^k))$. Then $R_{jq}^{ip} P_{pl}^{qk} = \delta_j^i \delta_l^k$ and P is the matrix of τ_{M,M^*} : $\tau_{M,M^*}(x_i \otimes \xi^j) = \xi^l \otimes x_k P_{il}^{jk}$. Further, set $C_j^i = u(a_j^i)$, $D_j^i = u(S(a_j^i))$. From the definition of u , we have $C_j^i = P_{jl}^{il}$, $D_j^i = P_{lj}^{ii}$. Equation (2) implies $CD = DC = T^{-2}$.

Assume that a morphism f has a matrix F . Then

$$\Phi(f) = \text{tr}(DTFA), \quad \Phi(f^*) = \text{tr}(CTFA), \quad (6)$$

where tr denotes the usual trace.

We define the braided power of the multiplicative matrix A to be the image of $\xi^i \otimes x_j$ under $\Phi_{M^{\otimes n-1}}(\tau_{M^{\otimes n-1}, M})$. Using the third property of θ , we have

$$(A^{q*n})_j^i = \Phi_{M^{\otimes n-1}}(\tau_{M^{\otimes n-1}, M})(\xi^i \otimes x_j) = (D^{\otimes n-1} T^{\otimes n-1})_J^I (R_{n-1} R_{n-2} \cdots R_1)_{Kk}^{Jj} A_I^K a_i^k, \quad (7)$$

where $a_J^I := a_{j_1}^{i_1} a_{j_2}^{i_2} \cdots a_{j_n}^{i_n}$.

1.3. Characters of Cosemisimple Coribbon Hopf Algebras. A Hopf algebra is called (right) cosemisimple iff the category of its (right) comodules is semisimple, that is all comodules are absolutely decomposable. In this case there exists an H -comodule morphism, called integral, $\int : H \longrightarrow \mathbb{K}$, that is, $\int$ satisfies the following relation: $\int(a) = a_{(1)} \int(a_{(2)})$, $\int(1) = 1$. H decomposes into the direct sum of its simple subcoalgebra $H = \oplus_{i \in I} H_i$, $H_0 \cong \mathbb{K}$ and $\int|_{H_i} = 0$ unless $i = 0$. As in the case of finite group characters, we define a scalar product on the set of characters in H :

$$\langle \Phi(M), \Phi(N) \rangle := \int(\Phi(M \otimes N^*)) = \int(\Phi(M)\Phi(N^*)). \quad (8)$$

From the preceding discussion, one sees that for a simple H -comodule M , $\int(\Phi(M)) = 0$ unless $M \cong \mathbb{K}$. Thus, we have the orthogonal relations for simple H -comodules

$$\langle \Phi(M), \Phi(N) \rangle = \begin{cases} 1 & \text{if } M \cong N \\ 0 & \text{if } M \not\cong N. \end{cases} \quad (9)$$

The direct decomposition of H into its simple subcoalgebras also implies that non-isomorphic simple comodules have different characters, and the set of characters are linear independent.

Proposition 2. *Let H be a co-semi-simple Hopf algebra. The subspace of H spanned by characters is a subalgebra of H and is isomorphic to the Grothendieck ring (defined over $\mathbb{K}$) of finite dimensional H -comodules.*

2. MATRIX QUANTUM GROUPS OF TYPE A_n AND THEIR CHARACTERS

Let V be a vector space of finite dimension over $\mathbb{K}$ with a fixed basis $x_1, x_2, \dots, x_d$. Then an operator $R : V \otimes V \longrightarrow V \otimes V$ can be given by its matrix R_{kl}^{ij} . R is called Hecke symmetry if it satisfies the Yang-Bater equation $R_1 R_2 R_1 = R_2 R_1 R_2$, $R_1 := R \otimes \text{id}_V$, $R_2 := \text{id}_V \otimes R$, the Hecke equation $(R - q)(R + 1) = 0$ and is closed, i.e., there exists a matrix P_{kl}^{ij} , such that $P_{jq}^{ip} R_{pj}^{qk} = \delta_l^i \delta_j^k$. We shall always assume that $q^n \neq 1, \forall n > 1$.

The name Hecke symmetry comes from the fact that such an operator induces a representation of the Hecke algebra $\mathcal{H}_n = \mathcal{H}_{q,n}$ on the power $V^{\otimes n}$ of V : $\rho_n(T_i) = R_i := \text{id}_V^{\otimes i-1} \otimes R \otimes \text{id}_V^{\otimes n-i-1}$. We denote, for convenience, $R_w := \rho_n(T_w)$. By means of ρ_n we shall some times identify an element of $\mathcal{H}_n$ with an endomorphism of $\text{End}(V^{\otimes n})$.

The Hecke symmetry R is called even Hecke symmetry of rank r if the anti-symmetrizer operator $Y_n := \sum_{w \in \mathfrak{S}_n} (-q)^{-l(w)} R_w$ has non-zero rank for $n = r$ and vanishes for $n = r + 1$.

Given a Hecke symmetry R , we define a bialgebra E_R and a Hopf algebra H_R in the following way. Let $\{z_j^i, t_j^i | i, j = 1, 2, \dots, d\}$ be a set of variables. $E_R := \mathbb{K} \langle \{z_j^i | i, j = 1, 2, \dots, d\} \rangle / (R_{mn}^{ij} z_k^m z_l^n = z_p^i z_q^j R_{kl}^{pq})$, $H_R := \mathbb{K} \langle \{z_j^i, t_j^i | i, j = 1, 2, \dots, d\} \rangle / (R_{mn}^{ij} z_k^m z_l^n = z_p^i z_q^j R_{kl}^{pq}, t_k^i z_j^k = z_k^i t_j^k = \delta_j^i)$. The coproduct on E_R and H_R is given by $\Delta(z_j^i) = z_k^i \otimes z_j^k$, $\Delta(t_j^i) = t_j^k \otimes t_k^i$. The antipode on H_R is given by $S(z_j^i) = t_j^i$, $S(t_j^i) = C_m^i z_n^m C^{-1}{}_{j,n}$, $C_j^i := P_{jl}^{il}$.

E_R and H_R are coquasitriangular bialgebra with the CQT structure given by $r(z_j^i, z_l^k) = R_{jl}^{ki}$. H_R is a coribbon Hopf algebra with the coribbon structure $t(z_j^i) = q^{(r+1)/2} \delta_j^i$.

E_R and H_R coact on V by the coaction $\delta(x_i) = x_j \otimes z_i^j$. Hence $V^{\otimes n}$ is a comodule of E_R and H_R , its dual is comodule of H_R . The coaction of E_R and the action of $\mathcal{H}_n$ on $V^{\otimes n}$ are centralizers of each other in $\text{End}_{\mathbb{K}}(V^{\otimes n})$. Thus, $\rho_n(\mathcal{H}_n) \cong \text{End}^{E_R}(V^{\otimes n})$ and simple E_R comodules have form $\text{Im}(\rho_n(E_\lambda))$, where E_λ is a primitive idempotent of $\mathcal{H}_n$. The natural map $E_R \longrightarrow H_R$ is injective, hence simple E_R -comodule are simple over H_R , too.

If R is an even Hecke symmetry of rank r then H_R is cosemisimple, its simple comodules are parameterized by $\mathbb{Z}$ -partitions, that is, the sequences of non-increasing integers of length r : $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$, $\lambda_i \in \mathbb{Z}$, $\lambda_i \geq \lambda_{i+1}$. Let M_λ denote the simple comodule corresponding to λ . In particular, $V \cong M_{(1)}$. The coefficients $c_{\mu\nu}^\lambda$ in the decomposition

$$M_\mu \otimes M_\nu \cong \bigoplus_{\lambda} c_{\mu\nu}^\lambda M_\lambda \quad (10)$$

are the same as for those of irreducible rational $\text{GL}(r)$ -representations, i.e., the Littlewood-Richardson coefficients. Further, let $-\lambda := (-\lambda_r, -\lambda_{r-1}, \dots, -\lambda_1)$, $\lambda^+ := (\lambda_1 - \lambda_r, \lambda_2 - \lambda_r, \dots, 0)$. Then

$$(M_\lambda)^* \cong M_{-\lambda}, \quad M_\lambda \cong M_{\lambda^+} \otimes M(1^r)^{\otimes \lambda_r}.$$

Notice that $M_{(1^n)}$ is isomorphic to the image of Y_n in $V^{\otimes n}$. It is shown that Y_r has rank 1 over $\mathbb{K}$ hence induces a group-like element D in E_R , called quantum determinant.

The interested reader is referred to [4, 5] for more details.

2.1. Characters of Comodules over H_R . Now, using the construction of Section 1 we can define characters of simple H_R -comodules. Notice that the matrix C introduced above is precisely $C_j^i = u(z_j^i)$. We also set $D_j^i = u(t_j^i) = P_{jl}^{li}$. Then, according to (2), $CD = DC = q^{-(r+1)}$.

Let $\lambda \vdash n$. Then the simple comodule M_λ is isomorphic to $\text{Im}(\rho_n(E_\lambda))$, where E_λ is a primitive idempotent corresponding to λ . Let us use the same character (E_λ) to denote the endomorphism on $V^{\otimes n}$ induced by E_λ . Then we have

Lemma 3.

$$\begin{aligned}\Phi(M_\lambda) &= q^{n(r+1)/2} \text{tr}(D^{\otimes n} E_\lambda Z^{\otimes n}) & \Phi(M_\lambda^*) &= q^{n(r+1)/2} \text{tr}(C^{\otimes n} E_\lambda \overline{T^{\otimes n}}) \\ \Phi(F) &= q^{n(r+1)/2} \text{tr}(D^{\otimes n} F Z^{\otimes n}) & \Phi(F^*) &= q^{n(r+1)/2} \text{tr}(C^{\otimes n} F \overline{T^{\otimes n}}),\end{aligned}$$

where F is the matrix of a morphism $f : V^{\otimes n} \longrightarrow V^{\otimes n}$, $\overline{T^{\otimes n}}^I_J = T^{\otimes n I'}_{J'}$.

Proof. The above formula follow immediately from the identity (4). Recall that Φ is a trace map from $\text{End}^{E_R}(V^{\otimes n})$ into H_R . Combining Φ with ρ_n we get a trace map $\mathcal{H}_n \longrightarrow H_R$. Using Lemma 1 we can represent the character of a comodule morphism on $V^{\otimes n}$ as a linear combination of $\Phi(T_{c_k})$, where $c_k := (1, 2, \dots, k)$ a k -cycle in $\mathfrak{S}_n$ (see [4]). Let $P_n := \Phi(T_{c_n})$, $H_n := \Phi(M_{(n)})$, $E_n := \Phi(M_{(1^n)})$.

Lemma 4. *We have the following relations*

$$\begin{aligned}[n]_q H_n &= H_{n-1} P_1 + H_{n-2} P_2 + \cdots + H_0 P_n, \\ [n]_{1/q} E_n &= E_{n-1} P_1 - q^{-1} E_{n-2} P_2 + \cdots + (-q)^{n-1} E_0 P_n,\end{aligned}$$

where $[n]_q := (q^n - 1)/(q - 1)$.

The second relation was established in [7] for slightly differently defined P_n 's.

Proof. Let X_n and Y_n be the quantum symmetrizer and anti-symmetrizer operators on $V^{\otimes n}$. They can be defined by induction:

$$\begin{aligned}[n]_q X_n &= (1 + R_{n-1} + \cdots + R_1 R_2 \cdots R_{n-1}) X_{n-1}, \\ [n]_{1/q} Y_n &= (1 - q^{-1} R_{n-1} + \cdots + (-q)^{n-1} R_1 T_2 \cdots R_{n-1}) Y_{n-1}.\end{aligned}$$

X_n and Y_n are projectors on $V^{\otimes n}$ and their images are isomorphic to $M_{(n)}$ and $M_{(1^n)}$ respectively. Hence $\Phi(X_n) = S_{(n)} = H_n$ and $\Phi(Y_n) = S_{(1^n)} = E_n$. We have $X_n R_i = R_i X_n = q X_n$, $Y_n R_i = R_i Y_n = -Y_n$. Therefore

$$\begin{aligned}[n]_q H_n &= \Phi((1 + R_{n-1} + \cdots + R_1 R_2 \cdots R_{n-1}) X_{n-1}) \\ &= \Phi(X_{n-1} (1 + R_{n-1} + \cdots + R_1 R_2 \cdots R_{n-1})) \\ &= H_{n-1} P_1 + \Phi((1 + R_{n-2} + \cdots + R_1 R_2 \cdots R_{n-2}) X_{n-2} R_{n-1}) \\ &= H_{n-1} P_1 + \Phi(X_{n-2} (1 + R_{n-2} + \cdots + R_1 R_2 \cdots R_{n-2}) R_{n-1}) \\ &= \cdots \\ &= H_{n-1} P_1 + H_{n-2} P_2 + \cdots + P_n.\end{aligned}$$

The second equation is proved analogously. ■

Corollary 5. *The character map Φ is a trace map on $\mathcal{H}_n$ with value in Λ_R , the subalgebra of E_R spanned by $\{S_\lambda := \Phi(M_\lambda) | \lambda \in \mathcal{P}^r\}$.*

On the other hand, according to Equation (9) and Lemma 1, we have

$$S_\lambda = \det |(H_{\lambda_i - i + j})_{1 \leq i, j \leq r}|, \lambda \vdash n \quad S_\lambda = S_{\lambda^+} \cdot D^{\lambda^r}. \quad (11)$$

Therefore we can define an algebra isomorphism $\Lambda_E \longrightarrow \Lambda_r$, $H_k \longmapsto h_k$, where Λ_r is the algebra of symmetric functions in r variables and h_k are k -th complete symmetric function. This isomorphism is in fact an isometry with respect to the scalar product defined in (6), under which S_λ is mapped to s_λ – the Schur functions and, in particular, $E_k := S_{(1^k)}$ is mapped to the elementary symmetric function e_k . Thus S_λ can be consider as quantum symmetric functions on non-commuting variables. Notice that the elements P_k are mapped to the symmetric functions q_k , introduced by A. Ram in [8].

Let K be an algebra over $\mathbb{K}$. A K -point $\mathcal{A}$ of E_R is an algebra homomorphism $\mathcal{A} : E_R \rightarrow K$. Let $A = \mathcal{A}(Z)$. Then $\mathcal{A}$ satisfies the relation $RA_1A_2 = A_1A_2R$. Conversely, each matrix of elements from K , satisfying this relation give a K -point of E_R , it is then called a quantum matrix with coefficients from K . In this case we can define the elements $S_\lambda(A)$ of K , replacing Z by A in the definition of S_λ . $S_\lambda(A)$ is called the value of S_λ at the point $\mathcal{A}$. Analogously, let E_R^{op} be the subbialgebra of H_R , generated by t 's. Let N be a point of E_R^{op} , we define $S_{-\lambda}(N)$ to be $S_{-\lambda}$ computed at $T = N$.

Let us consider an example. Assume that R is the Drinfel'd-Jimbo matrix of type A_{r-1} , for $1 \leq i, j \leq r$, $p^2 = q$,

$$R_{ij}^{kl} := \begin{cases} p^2 & \text{if } i = j = k = l \\ p^2 - 1 & \text{if } k = i < j = l \\ p & \text{if } k = j \neq i = l \\ 0 & \text{otherwise} \end{cases}$$

Then R is an even Hecke symmetry of rank r . The associated Hopf algebra H_R is called (the function algebra on) quantum general linear group $GL_q(r)$. The elementary symmetric functions are

$$E_k = p^{k(r+1)} \sum_{i_1 < i_2 < \dots < i_k} p^{2(i_1 + i_2 + \dots + i_k)} \sum_{\sigma \in \mathfrak{S}_k} (-p)^{-l(\sigma)} z_{i_1}^{i_{1\sigma}} z_{i_2}^{i_{2\sigma}} \dots z_{i_k}^{i_{k\sigma}}.$$

We see that $R_{ij}^{kl} = 0$ unless $(i, j) = (k, l)$ or $(i, j) = (l, k)$. Therefore any diagonal matrix with commuting entries in the diagonal is a point of $GL_q(r)$. Let $A = (a_1, a_2, \dots, a_r)$ be such a point. Then

$$S_\lambda(A) = s_\lambda(qa_1, q^2a_2, \dots, q^ra_r),$$

where s_λ are the usual Schur functions.

2.2. The Characteristic Equation. Using the construction of Section 1 we can also define the quantum powers of the matrix Z .

$$(Z^{q*n})_j^i := q^{n(r+1)/2} D_N^{\otimes n M} (R_n R_{n-1} \dots R_1)_{Pq}^{Nj} Z_M^P z_i^q$$

Theorem 6. (The characteristic equation for quantum matrix) The quantum matrix Z satisfies the following equation

$$Z^{q*r} - E_1 Z^{q*(r-1)} + \dots + (-1)^r E_r = 0. \quad (12)$$

Proof. If R is an even Hecke symmetry of rank r , $Y_{r+1} = 0$. Taking the partial trace of Y_{r+1} with respect to $V^{\otimes r}$, we have

$$0 = \Phi_{V^{\otimes r}}((1 - q^{-1}R_r + \dots + (-q)^{-r}R_1R_2 \dots R_r)Y_r)$$

Using the equation (5), we have

$$\begin{aligned} & q^{n(r+1)/2} \Phi_{V^{\otimes r}}((1 - q^{-1}R_r + \dots + (-q)^{-r}R_1R_2 \dots R_r)Y_r) \\ &= \Phi_{V^{\otimes r}}(Y_r(1 - q^{-1}R_r + \dots + (-q)^{-r}R_1R_2 \dots R_r)) \\ &= \Phi_{V^{\otimes r}}(Y_r) - [r]_{1/q} \Phi_{V^{\otimes r}}(Y_r R_r) \\ &= q^{-n(r+1)/2} E_r \cdot Z - \Phi_{V^{\otimes r}}((1 - q^{-1}R_{r-1} + \dots + (-q)^{1-r}R_1R_2 \dots R_{r-1})Y_{r-1}R_r) \\ &= \dots \\ &= q^{-n(r+1)/2} (E_r \cdot Z - E_{r-1} \cdot Z^{q*2} + \dots + (-1)^r Z^{q*(r+1)}). \end{aligned}$$

Thus,

$$E_r \cdot Z - E_{r-1} \cdot Z^{q*2} + \dots + (-1)^r Z^{q*(r+1)} = 0. \quad (13)$$

We have the following equalities

$$\begin{aligned} (Z^{q*n})_i^j t_m^i P_{jn}^{mk} D^{-1}_l^n &= q^{(r+1)/2} (Z^{q*(n-1)})_l^k, \\ z_i^j t_m^i P_{jn}^{mk} D^{-1}_l^n &= q^{(r+1)/2} \delta_l^k. \end{aligned}$$

Multiplying the left-hand side of (13) with $t_m^i P_{jn}^{mk} D^{-1}_l^n$, we obtain

$$Z^{q*r} - E_1 Z^{q*(r-1)} + \dots + (-1)^r E_r = 0.$$

A priori this equation holds in H_R . However, since the canonical map $E_R \rightarrow H_R$ is injective, this equation also holds in E_R . ■

3. THE q -ANALOGUE OF HARISH-CHANDRA-ITZYKSON-ZUBER INTEGRAL

The HCIZ integral on $U(n)$, $\int e^{\text{tr}(MUNU^\dagger)} dU$, for any Hermitian matrices M and N , is given by

$$\int e^{\text{tr}(MUNU^\dagger)} dU = \sum_{n=1}^{\infty} \frac{t^n}{n!} \sum_{\lambda \vdash n} \frac{d_\lambda}{r_\lambda} \Phi_\lambda(M) \Phi_\lambda(N).$$

Here d_λ is the dimension of the irreducible representation S_λ of the symmetric group $\mathfrak{S}_n$ and r_λ is the dimension of the irreducible representation M_λ of $U(n)$, Φ_λ is the character of M_λ , thus $\Phi_\lambda(M)$ is the symmetric function s_λ computed at the eigenvalues of M . Equivalently, we have

$$\int \text{tr}(MUNU^\dagger)^n dU = \sum_{\lambda \vdash n} \frac{d_\lambda}{r_\lambda} \Phi_\lambda(M) \Phi_\lambda(N).$$

In this section we want to give an analogue of this formula for the integral on H_R .

A formula for the normalized integral on H_R is given in [5]. Since H_R is coquasi-triangular, its elements can be represented as linear combinations of elements of the form $z_{i_1}^{j_1} z_{i_2}^{j_2} \dots z_{i_n}^{j_n} t_{k_1}^{l_1} t_{k_2}^{l_2} \dots t_{k_m}^{l_m}$. For simplicity we denote $Z_I^J := z_{i_1}^{j_1} z_{i_2}^{j_2} \dots z_{i_n}^{j_n}$ and so on. The integral on $Z_I^J T_K^L$ vanishes unless $l(I) = l(K)$, where $l(I)$ means the length of I . If $l(I) = l(K) = n$, then

$$\int Z_I^J T_K^L = \sum_{w \in \mathfrak{S}_n} (P_n C^{\otimes n} R_{w^{-1}})_I^{L'} T_{wK'}^J \quad (14)$$

where, for $K = (k_1, k_2, \dots, k_n)$, $K' := (k_n, k_{n-1}, \dots, k_1)$.

$$P_n := \prod_{k=1}^n (L_k - [-r]_q)^{-1}, \quad L_k := \sum_{i=1}^{n-1} q^{-i} R_{(n-i,i)}; \quad L_1 := 0.$$

L_k is the Murphy operator introduced in [2]. Let F_λ be the minimal central idempotent in $\mathcal{H}_n$, corresponding to $\lambda \vdash n$. Then

$$P_n = \sum_{\lambda \vdash n} q^{r_n} \prod_{x \in [\lambda]} [c_\lambda(x) + r]_q^{-1} F_\lambda,$$

$c_\lambda(x)$ is the content of the node x in the diagram $[\lambda]$ associated to λ , if x is in the i^{th} row and j^{th} column then $c_\lambda(x) = j - i$. Let us denote $p_\lambda := q^{r_n} \prod_{x \in [\lambda]} [c_\lambda(x) + r]_q^{-1}$.

$\mathcal{H}_n$ possesses a non-degenerate symmetric associative bilinear form, given by $\langle T_u, T_w \rangle := \delta_u^{v^{-1}} q^{l(u)}$. Then $\chi(h) = \langle h, 1 \rangle$ is a trace map on $\mathcal{H}_n$. By definition, $\chi(h)$ is the coefficient of 1 in the presentation of h as the linear combination of T_w . We wish to find $\chi(E_\lambda)$.

Let $\mathcal{H}_n = \bigoplus_{\lambda \in \mathcal{P}_n} \mathcal{A}_\lambda$ be the decomposition of $\mathcal{H}_n$ into the direct sum of its minimal two-sided ideals. $\mathcal{A}_\lambda$ is itself a matrix ring of degree d_λ , with the unit F_λ . Let E_λ^{ij} be a $\mathbb{K}$ basis of $\mathcal{A}_\lambda$ such that $E_\lambda^{ij} E_\lambda^{kl} = \delta_k^j E_\lambda^{il}$. From standard argument (cf. [3]) for any trace map h on $\mathcal{H}_n$, $h(E_\lambda^{ij}) = 0$ unless $i = j$ and $h(E_\lambda^{ii}) = d_\lambda^{-1} \chi(F_\lambda) =: k_\lambda$ does depend only on λ . Then $\langle E_\lambda^{ij}, E_\mu^{kl} \rangle = \delta_\lambda^\mu \delta_j^k \delta_i^l k_\lambda$. Therefore $\{\frac{1}{k_\lambda} E_\lambda^{ji} | \lambda \in$

$\mathcal{P}_n, i, j = 1, 2, \dots, d_\lambda\}$ is the dual basis to $\{E_\lambda^{ij} | \lambda \in \mathcal{P}_n, i, j = 1, 2, \dots, d_\lambda\}$ with respect to the bilinear form defined above. On the other hand, with respect to this bilinear form, $\{q^{-l(w)}T_{w^{-1}} | w \in \mathfrak{S}_n\}$ is the dual basis to $\{T_w | w \in \mathfrak{S}_n\}$. Thus, we have

$$\sum_{w \in \mathfrak{S}_n} q^{-l(w)}T_{w^{-1}} \otimes T_w = \sum_{\lambda, i, j} \frac{1}{k_\lambda} E_\lambda^{ij} \otimes E_\lambda^{ji}. \quad (15)$$

Now assume that M is a point of E_R and N is a point of E_R^{op} . According to (14), (15), we have

$$\begin{aligned} & \int q^{n(r+1)/2} \text{tr}(D^{\otimes n} M^{\otimes n} Z^{\otimes n} \overline{N^{\otimes n} T^{\otimes n}}) \\ &= \sum_{w \in \mathfrak{S}_n} q^{-l(w)} q^{n(r+1)/2} \text{tr}(P_n C^{\otimes n} R_{w^{-1}} \overline{N^{\otimes n}}) \cdot \text{tr}(R_w D^{\otimes n} M^{\otimes n}) \\ &= \sum_{\substack{1 \leq i, j \leq d_\lambda \\ \lambda \vdash n}} k_\lambda^{-1} q^{n(r+1)/2} \text{tr}(P_n E_\lambda^{ij} C^{\otimes n} \overline{N^{\otimes n}}) \cdot \text{tr}(E_\lambda^{ij} D^{\otimes n} M^{\otimes n}) \\ &= \sum_{\substack{1 \leq i, j \leq d_\lambda \\ \lambda \vdash n}} k_\lambda^{-1} p_\lambda q^{-n(r+1)/2} \Phi(E_\lambda^{ij*})(N) \cdot \Phi(E_\lambda^{ji})(M) \\ &= \sum_{\lambda \vdash n} q^{-n(r+1)/2} d_\lambda k_\lambda^{-1} p_\lambda S_\lambda(M) S_{-\lambda}(N). \end{aligned}$$

Here $\overline{N^{\otimes n}}^I_J := M^{\otimes n}_{J'}$. In the last equation we use the fact that $\Phi(E_\lambda^{ji}) = 0$ unless $i = j$. Thus, to compute the integral above, it remains to find k_λ .

Since $\mathcal{H}_n$ is semi-simple, Φ can be uniquely represented as a combination of irreducible characters of $\mathcal{H}_n$ with coefficients from Λ_E : $\Phi = \sum_\lambda K_\lambda \chi^\lambda$, where χ^λ is the irreducible character of $\mathcal{H}_n$, that corresponds to $\lambda \in \mathcal{P}_n$. Let E_λ be a primitive idempotent corresponding to λ , then $\chi^\lambda(E_\mu) = \delta_\mu^\lambda$ and $\Phi(E_\lambda) = S_\lambda$, hence $K_\lambda = S_\lambda$. Thus, we have $\Phi = \sum_\lambda \chi^\lambda S_\lambda$. The orthogonal relations in (6) imply

$$\chi^\lambda(W) = (\Phi(W), \Phi(E_\lambda)) = \int (\Phi(W) \Phi(E_\lambda^*)).$$

Recall that (cf. (6)) $\Phi(W) = q^{n(r+1)/2} (D^{\otimes n} W Z^{\otimes n})$, $\Phi(W^*) = q^{n(r+1)/2} (C^{\otimes n} W Z^{\otimes n})$. Therefore using (14), (15), we have

$$\begin{aligned} \chi^\lambda(W) &= \sum_{w \in \mathfrak{S}_n} q^{-l(w)} \text{tr}(C^{\otimes n} E_\lambda^{ii} T_w W P_n T_{w^{-1}}) \\ &= k_\lambda^{-1} p_\lambda \text{tr}(C^{\otimes n} W F_\lambda), \end{aligned}$$

here we use the fact that $CD = q^{-(r+1)}$. In particular, $1 = \chi^\lambda(E_\lambda^{ii}) = k_\lambda^{-1} p_\lambda \text{tr}(C^{\otimes n} E_\lambda^{ii})$.

Notice that $q^{r(n+1)/2} \text{tr}(C^{\otimes n} W) = \varepsilon(\Phi(W^*))$ is the braided trace of the morphism W in $\text{End}(V^{\otimes n})$. In particular, $q^{r(n+1)/2} \text{tr}(C^{\otimes n} E_\lambda^{ii})$ is the rank of the simple comodule M_λ , it was computed in [5]:

$$\text{rank}(M_\lambda) = q^{-n(r-1)/2+n(\lambda)} \prod_{x \in [\lambda]} \frac{[c_\lambda(x) + r]_q}{[h_\lambda(x)]_q}.$$

Therefore

$$k_\lambda = q^{n(\lambda)} \prod_{x \in [\lambda]} [h(x)]_q^{-1}.$$

Thus we have proved

Proposition 7. *The value of the character χ on a primitive idempotent E_λ is equal to $q^{n(\lambda)} \prod_{x \in [\lambda]} [h(x)]_q^{-1}$, or equivalently $\chi = \sum_\lambda q^{n(\lambda)} \prod_{x \in [\lambda]} [h(x)]_q^{-1} \chi^\lambda$.*

The value of k_λ gives us immediately the quantum analogue of HCIZ integral.

Theorem 8. *The quantum HCIZ integral on quantum group of type A_{r-1} is given by*

$$\int q^{n(r+1)/2} \text{tr}(D^{\otimes n} M^{\otimes n} Z^{\otimes n} \overline{N^{\otimes n} T^{\otimes n}}) = \sum_{\lambda \vdash n} \frac{d_\lambda}{\text{rank}(M_\lambda)} S_\lambda(M) S_{-\lambda}(N).$$

Example. Assume that R is the Drinfel'd-Jimbo matrix of type A_{r-1} and $M = (\mu_1, \mu_2, \dots, \mu_r)$, $N = (\nu_1, \nu_2, \dots, \nu_r)$ be diagonal matrices. Then we have

$$\int (q^{n(r+1)/2} q^{|I|} \mu^I Z_J^I \nu^J T_I^J) = \sum_{\lambda \vdash n} \frac{d_\lambda}{\text{rank}(M_\lambda)} s_\lambda(q^i \mu_i) s_\lambda(q^i \nu_i),$$

where $|I| := i_1 + i_2 + \dots + i_m$, $\mu^I := \mu^{i_1} \mu^{i_2} \dots \mu^{i_m}$.

Notice that the value of k_λ allows us to give a formula for computing irreducible characters of the Hecke algebra $\mathcal{H}_n$. In fact, since χ is faithful, an irreducible character χ^λ can be computed from χ in the following sense $\chi^\lambda(W) = \chi(W F_\lambda) \chi(F_\lambda)^{-1}$, where F_λ is the minimal central idempotent corresponding to λ . If $\text{rank}(R) = n$ then $\mathcal{H}_n \cong \text{End}^{E_R}(V^{\otimes n})$. Therefore we have

Proposition 9. *The minimal central idempotents F_λ contain all information about the character table of the Hecke algebra. Let $t_\lambda(w)$ be the coefficient of T_w in F_λ . Then*

$$\chi^\lambda(T_w) = \frac{q^{l(w)} q^{-n(\lambda)}}{n!} \prod_{x \in \lambda} (h(x) [h(x)]_q) t_\lambda(w^{-1}).$$

Indeed, $\chi(F_\lambda) = d_\lambda \chi(E_\lambda) = q^{n(\lambda)} d_\lambda \prod_{x \in [\lambda]} [h(x)]_q^{-1}$. $d_\lambda = n! \prod_{x \in [\lambda]} h(x)^{-1}$.

Acknowledgment

The author would like to thank the Abdus Salam International Centre for Theoretical Physics and the Max-Planck Institute for Mathematik, Bonn, for their hospitality and financial supports.

REFERENCES

- [1] C. C. Itzykson and J.B. Zuber. The planar approximation. II. *J. Math. Phys.*, 21:411–421, 1980.
- [2] R. Dipper and G. James. Block and Idempotents of Hecke Algebras of General Linear Groups. *Proc. London Math. Soc.*, 54(3):57–82, 1987.
- [3] F.M. Goodman, P. de la Harpe, and V.F.R. Jones. *Coxeter Graphs and Towers of Algebras*, volume 14 of *MSRI publications*. Springer-Verlag, 1989.
- [4] Phung Ho Hai. Koszul Property and Poincaré Series of Matrix Bialgebra of Type A_n . *Journal of Algebra*, 192(2):734–748, 1997.
- [5] Phung Ho Hai. On Matrix Quantum Groups of Type A_n . *Preprint ICTP/97101, q-alg/9708007*, 1997.
- [6] A. Joyal and R. Street. Braided Tensor Categories. *Advances in Math.*, 102:20–78, 1993.
- [7] P.N. Pyatov and P.A. Saponov. Newton Relations for Quantum Matrix Algebras of RTT -Type. *Preprint IHEP/96-76*, 1996.
- [8] A. Ram. A Frobenius Formula for The Characters of The Hecke Algebras. *Invent. of Math.*, 106:461–468, 1991.
- [9] N. Reshetikhin and V. Turaev. Ribbon Graph and Their Invariant Derived from Quantum Groups. *Comm. Math. Phys.*, 127:1–26, 1990.

HANOI INSTITUTE OF MATHEMATICS P.O. BOX 631, 10000 BO HO, HANOI, VIETNAM

Current address: Max-Planck Institut für Mathematik, Gottfried-Claren-Str. 26, 53225, Bonn, Germany

E-mail address: phung@mpim-bonn.mpg.de